

UNCLASSIFIED

ON PERTURBATIONS OF STABILITY INEQUALITIES WITH AN
APPLICATION TO FINITE... (U) WISCONSIN UNIV-MADISON
MATHEMATICS RESEARCH CENTER R D GRIGORIEFF SEP 83
MRC-TSR-2564 DAAG29-80-C-0041 F/G 12/1


178

NL

END

DATE
CUMED

83
DTIC



MICROCOPY RESOLUTION TEST CHART
 NA - 1010-A

MRC Technical Summary Report #2564

ON PERTURBATIONS OF STABILITY INEQUALITIES
WITH AN APPLICATION TO FINITE DIFFERENCE
APPROXIMATIONS OF ODE'S

R. D. Grigorieff

AD-A194 535

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

September 1983

(Received March 7, 1983)

**Approved for public release
Distribution unlimited**

DTIC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

**DTIC
ELECTE**
NOV 9 1983

24
A

83 11 68 149

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

ON PERTURBATIONS OF STABILITY INEQUALITIES WITH AN APPLICATION
TO FINITE DIFFERENCE APPROXIMATIONS OF ODE'S

R. D. Grigorieff*

Technical Summary Report #2564
September 1983

ABSTRACT

↓
In the framework of Stummel's discrete approximation theory, a perturbation theorem for inverse stability inequalities is proven. As an application, the inverse stability of compact finite difference schemes approximating two-point boundary value problems for linear ordinary differential equations on nonuniform grids is established. K

AMS (MOS) Subject Classifications: 65J10, 65L07, 65L10

Key Words: Discrete approximation, inverse stability, ODE, BVP,
finite differences, nonuniform grids

Work Unit Number 3 - Numerical Analysis and Scientific Computing

*Technical University of Berlin, Str. des 17. Juni 135, D-1000 Berlin 12,
West Germany

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.



SIGNIFICANCE AND EXPLANATION

The normal procedure in solving a continuously defined problem numerically consists in applying a discretization first which reduces the original problem to one which can be treated by numerical algorithms for solving equation with a finite number of unknowns. A fundamental question arising in this context is the convergence of the solutions of the discretized problem to the solution of the original one.

In studying these kinds of questions, it has turned out that the various different discretization procedures can be treated in a unified manner in the abstract setting of the so-called discrete approximation theory which is also used in this paper. The main point in proving convergence in the stability of the discretization. This paper deals with methods in proving stability for linear problems. As an application of the abstract result obtained, the stability of finite difference approximations for linear two-point boundary value problem on not equally spaced grids is established, a topic which has attracted the attention of a number of numerical analysts in the last few years.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

ON PERTURBATIONS OF STABILITY INEQUALITIES WITH AN APPLICATION
TO FINITE DIFFERENCE APPROXIMATIONS OF ODE'S

R. D. Grigorieff*

1. Introduction. The main part in proving the convergence of finite difference approximations for boundary value problems consists in establishing suitable stability inequalities. One common method in proving such inequalities is to start from an often comparatively simple a-priori-inequality which leads to the desired stability by a perturbation argument. The purpose of this report is to give an abstract version of this procedure in the framework of the discrete approximation theory introduced by Stummel [10,11]. In this way it can be clearly recognized which properties of the problem give rise to the stability inequalities. As a further consequence one obtains fairly simple proofs of stability results used only very indirectly in the concrete context or being even not found there (e.g. [9,12]).

As an application we establish the inverse stability of compact finite difference approximations for linear m -th order two-point boundary value problems on nonuniform grids in the maximum norm, which makes it easy to get the convergence of the schemes obtained in [9,12]. These stability inequalities have also been given in [4] using a different manner of proof. The method developed in this report opens the possibility to treat also more general schemes as well along with the associated eigenvalue problem. Moreover the abstract result applies equally well to other discretization methods, e.g. to Galerkin and quadrature methods, a fact which is mentioned only briefly here but which is wellknown to those familiar with discrete approximations.

*Technical University of Berlin, Str. des 17. Juni 135, D-1000 Berlin 12, West Germany

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

2. Notations. Let a denumerable sequence Λ_0 and normed spaces $E, E_1, \dots, E_{\Lambda_0}$ be given. The spaces are said to form a discrete approximation $A(E, \{E_{\Lambda_1}\})$ if a linear map Lim is defined on a linear subset of all sequences $u_1 \in E_1, \dots, u_{\Lambda_0} \in E_{\Lambda_0}$, with range equal to E and the property

$$\text{Lim}(u_1, \dots, u_{\Lambda_0}) = 0 \iff \|u_1\| + 0 \quad (1 \in \Lambda_0)$$

(the same symbol is used here to denote all the various norms on E, E_1). A sequence $u_1, \dots, u_{\Lambda_0}$ lying in the domain of Lim is said to be discretely convergent and we write

$$u_1 \rightarrow u \quad (1 \in \Lambda_0)$$

if u is its image under the map Lim . By Λ_1, Λ_2 we denote final sections of

Λ_0 , by Λ, Λ' subsequences of Λ_0 , not necessarily the same at different occurrences.

The convergence of a subsequence u_1, \dots, u_{Λ} to $u \in E$ is defined in the obvious manner:

$$u_1 \rightarrow u \quad (1 \in \Lambda) \iff \forall v_1, \dots, v_{\Lambda_0} : v_1 \rightarrow u \quad (1 \in \Lambda_0), \|u_1 - v_1\| + 0 \quad (1 \in \Lambda).$$

By B_1 we denote the closed ball of radius 1 in E_1 . Let $G_1, \dots, G_{\Lambda_0}$ be subsets of E_1 . Then we introduce the limit set

$$\text{Lim sup } G_1 = \{u \in E \mid \exists \Lambda \in \Lambda_0, u_1 \in G_1, \dots, u_{\Lambda} \in G_{\Lambda}, u_1 \rightarrow u \quad (1 \in \Lambda)\}.$$

The sequence $G_1, \dots, G_{\Lambda_0}$ is said to be (locally) discretely compact, if each (bounded) sequence $u_1 \in G_1, \dots, u_{\Lambda} \in G_{\Lambda}$ contains a convergent subsequence.

Let also $A(F, \{F_{\Lambda_1}\})$ be a discrete approximation of normed spaces. A sequence of linear mappings $L_1 : E_1 \rightarrow F_1, \dots, L_{\Lambda_0} : E_{\Lambda_0} \rightarrow F_{\Lambda_0}$ is said to be discretely convergent to a linear mapping $L : E \rightarrow F$, in symbols $L_1 \rightarrow L \quad (1 \in \Lambda_0)$, if

$$u_1 \rightarrow u \quad (1 \in \Lambda_0) \implies L_1 u_1 \rightarrow L u \quad (1 \in \Lambda_0).$$

The sequence $L_1, \dots, L_{\Lambda_0}$ is said to be consistent with L if

$$\forall u \in E, u_1 \in E_1, \dots, u_{\Lambda} \in E_{\Lambda} : u_1 \rightarrow u, L_1 u_1 \rightarrow L u \quad (1 \in \Lambda_0).$$

The sequence $L_1, \dots, L_{\Lambda_0}$ is said to be discretely compact if the sequence

$R(L_1), \dots, R(L_{\Lambda_0})$ of ranges of L_1 is discretely compact. Sometimes we use the notation $E_1, \dots, E_{\{0\}}$, to indicate the space E , and similarly for F_1, L_1 etc.

3. The perturbation theorem. Let $A(E, \{E_{\Lambda_1}\}), A(F, \{F_{\Lambda_1}\}), A(G^{(j)}, \{G_{\Lambda_1}^{(j)}\}), j = 1, 2$, be discrete approximations of normed spaces. Let

$$L_1^{(j)} : E_1 \rightarrow F_1, M_1^{(j)} : E_1 \rightarrow G_1^{(j)}, i \in \Lambda_0 \cup \{0\}, j = 1, 2$$

be linear mappings. We wish to conclude that the stability inequality

$$(1) \quad \gamma_2 \|u_1\| \leq \|L_1^{(2)} u_1\| + \|M_1^{(2)} u_1\|, u_1 \in E_1, i \in \Lambda_2$$

holds if the inequality

$$(2) \quad \gamma_1 \|u_1\| \leq \|L_1^{(1)} u_1\| + \|M_1^{(1)} u_1\|, u_1 \in E_1, i \in \Lambda_1$$

is known to hold. Here γ_1, γ_2 denote positive constants independent of u_1 and i .

For brevity we set

$$K_1 : L_1^{(1)} - L_1^{(2)}, i \in \Lambda_0 \cup \{0\}.$$

(3) Let (2) and the following conditions be fulfilled:

(i) The sequence $K_1, i \in \Lambda_0$, is discretely compact and convergent to K

(ii) The sequence $M_1^{(1)}(B_1)$ is locally discretely compact

(iii) The sequence $(L_1^{(1)}, M_1^{(1)})$ is consistent with $(L^{(1)}, M^{(1)})$

(iv) $M_1^{(2)} + M^{(2)} (i \in \Lambda_0)$

(v) $(w, g) \in \text{Lim sup } (K_1(B_1), G^{(1)}) \cap \text{Lim sup } (L_1^{(1)}, M_1^{(1)})(B_1) \Rightarrow$

$$\exists u \in E : L^{(1)} u = w, M^{(1)} u = g$$

(vi) $L^{(2)} u = 0, M^{(2)} u = 0 \Rightarrow u = 0.$

Then inequality (1) holds.

Proof. Suppose (1) not to hold. Then there exist a subsequence $\Lambda \subset \Lambda_0$ and elements

$u_1 \in E_1, i \in \Lambda$, such that

$$(4) \quad \|u_1\| = 1, L_1^{(2)} u_1 \rightarrow 0, M_1^{(2)} u_1 \rightarrow 0 (i \in \Lambda).$$

Because of the assumption (i), we choose Λ such that also

$$K_1 u_1 \rightarrow w (i \in \Lambda)$$

for some $w \in F$. We now distinguish two cases.

First case: The sequence $M_1^{(1)} u_1, i \in \Lambda$, is bounded. Then, since (ii) is assumed,

without loss of generality

$$M_1^{(1)} u_1 \rightarrow g (i \in \Lambda)$$

for some $g \in G^{(1)}$. With (w, g) given in this way we can find a u with

$$(5) \quad L^{(1)} u = w, M^{(1)} u = g.$$

Now choose a consistency sequence $y_i, i \in \Lambda$, i.e.

$$y_i \rightarrow u, L_i^{(1)} y_i \rightarrow L^{(1)} u, M_i^{(1)} y_i \rightarrow M^{(1)} u \quad (i \in \Lambda).$$

Replacing u_i in (2) by $u_i - y_i$, one obtains

$$\gamma_i \|u_i - y_i\| < \|L_i^{(2)} u_i + K_i u_i - L_i^{(1)} u_i\| + \|M_i^{(1)} (u_i - y_i)\| \rightarrow 0 \quad (i \in \Lambda)$$

which shows the convergence $u_i \rightarrow u \quad (i \in \Lambda)$. From this, (4) and (i), (iv)

$$K_i u_i \rightarrow Ku = w, M_i^{(2)} u_i \rightarrow M^{(2)} u = 0 \quad (i \in \Lambda)$$

and so, because of (5), we conclude $L^{(2)} u = 0$, which gives $u = 0$. But as a consequence of (4) it is seen $u \neq 0$ and we have reached a contradiction.

Second case: The sequence $M_i^{(1)} u_i, i \in \Lambda$, is unbounded. By renorming u_i and if necessary passing to a subsequence which we continue to denote by Λ we obtain a sequence

$u_i \in E_i$ such that

$$(6) \quad \|u_i\| \rightarrow 0, L_i^{(2)} u_i \rightarrow 0, M_i^{(2)} u_i \rightarrow 0, \|M_i^{(1)} u_i\| = 1 \quad (i \in \Lambda).$$

We now proceed in the same way as in the first case, this time knowing $g \neq 0$. Since $g = M^{(1)} u$ and $u = 0$ this establishes the desired contradiction.

4. Special cases. The following specialization of theorem (3) is tailored to the treatment of an ordinary differential operator under two different sets of boundary conditions (see [1,2,4,7]). By $N(L)$ we denote the kernel of the mapping L .

(7) Let linear mappings

$$L_i : E_i \rightarrow F_i, M_i^{(j)} : E_i \rightarrow G, i \in \Lambda_0 \setminus \{0\}, j = 1, 2$$

be given such that with a constant $\gamma_1 > 0$

$$(8) \quad \gamma_1 \|u_i\| < \|L_i u_i\| + \|M_i^{(1)} u_i\|, u_i \in E_i, i \in \Lambda_1$$

holds and let the following conditions be satisfied:

$$(i) \quad \dim N(L) = \dim G < \infty$$

(ii) The restrictions $M^{(j)}|_{N(L)}, j = 1, 2$, are injective

(iii) The sequence $(L_i, M_i^{(1)}), i \in \Lambda_0$, is consistent with $(L, M^{(1)})$

$$(iv) \quad M_i^{(2)} \rightarrow M^{(2)} \quad (i \in \Lambda_0).$$

Then there exists $\gamma_2 > 0$ such that

$$(3) \quad \gamma_2 \|u_i\| < \|L_i u_i\| + \|M_i^{(2)} u_i\|, u_i \in E_i, i \in \Lambda_2.$$

Proof. We take $L_1^{(1)} = L_1^{(2)}$, $G_1 = G_1^{(1)} = G_1^{(2)} = G^{(1)} = G^{(2)} = G$, $K_1 = 0$, $K = 0$ in (3) with the discrete convergence in $A(G, \mathbb{R}G_1)$ to be the convergence in G . Then (3)(i)-(iv), (vi) hold. But also (3)(v) is satisfied since $w = 0$ and $M^{(1)}$ is surjective on $N(L)$ due to the assumptions (7)(i), (ii).

The next specialization of (3) is motivated by the wish to start from an a-priori inequality for the operator $L_1^{(1)}$, and so obtain a stability inequality for the operator $L_1^{(2)}$ differing from $L_1^{(1)}$ by "lower order terms".

(12) Let the following conditions be satisfied:

(i) $E_1 \subset G_1^{(1)}$, $1 \in \Lambda_0 \setminus \{0\}$, algebraically and topologically, and the sequence of natural imbeddings $J_1 : E_1 \rightarrow G_1^{(1)}$, $1 \in \Lambda_0$, is discretely compact and convergent to the natural imbedding $J : E \rightarrow G^{(1)}$

(ii) The sequence K_1 , $1 \in \Lambda_0$, is discretely compact and convergent to K

(iii) The sequence $L_1^{(1)}$, $1 \in \Lambda_0$, is consistent with $L^{(1)}$

(iv) $J_1 u_1 \rightarrow g$, $L_1^{(1)} u_1 \rightarrow w$ ($1 \in \Lambda$) $\Rightarrow u \in E : L^{(1)} u = w, Ju = g$

(v) $M_1^{(2)} + M^{(2)}$ ($1 \in \Lambda_0$)

(vi) $L^{(2)} u = 0$, $M^{(2)} u = 0 \Rightarrow u = 0$.

If the a-priori inequality

(13) $\gamma_1 \|u_1\| < \|L_1^{(1)} u_1\| + \|J_1 u_1\|$, $u_1 \in E_1$, $1 \in \Lambda_1$,

holds with some $\gamma_1 > 0$, then also (1) holds with some constant $\gamma_2 > 0$.

Proof. Conditions (3)(i), (iv)-(vi) directly correspond to (12)(ii), (iv)-(vi). Since $M_1^{(1)} = J_1$ and J_1 , $1 \in \Lambda_0$, is discretely compact, (3)(ii) is satisfied. The consistency (3)(iii) follows from (12)(iii) and the convergence $J_1 \rightarrow J$ assumed in (12)(i).

It should be remarked that the condition $Ju = g$ in (12)(iv) is only notational since it only distinguishes the element $u \in E$ and its imbedding into $G^{(1)}$.

The last specialization of theorem (3) we give is connected with the study of discretizations of equations of the second kind with a compact operator.

(14) Let the following conditions be satisfied:

(i) $\gamma_1 \|u_1\| < \|L_1 u_1\|$, $u_1 \in E_1$, $1 \in \Lambda_1$ with some $\gamma_1 > 0$

(ii) $L : E \rightarrow F$ is surjective

(iii) The sequence K_1 , $1 \in \Lambda_0$, is discretely compact and convergent to K

(iv) $(L-K)u = 0 \Rightarrow u = 0$

(v) The sequence L_1 , $1 \in \Lambda_0$, is consistent with L .

Then, for some $\gamma_2 > 0$,

$$\gamma_2 \|u_1\| < \|(L_1 - K_1)u_1\|, u_1 \in E_1, 1 \in \Lambda_2.$$

Proof. We apply theorem (3) with $L_1^{(1)} := L_1$, $M_1^{(1)} = M_1^{(2)} := 0$, $1 \in \Lambda_0 \setminus \{0\}$. It is easy to check that all assumptions of (3) are satisfied.

5. An application. In this section we wish to show how the convergence theorem contained in [9,12] can be derived from the results of this note. Incidentally, we establish a stability inequality which has been looked for in [12, p. 743].

In [9,12] compact, implicit difference schemes have been derived for a single m -th order differential equation

$$(16) \quad Lu(t) := u^{(m)}(t) + \sum_{i < m} a_i(t) u^{(i)}(t), t \in [A, B],$$

with boundary conditions

$$(17) \quad M^{(q)}u := \mu^{(q)}[u] + \nu^{(q)}[u] := \sum_{i=0}^m a_i^{(q)} u^{(i)}(A) + \sum_{i=0}^m b_i^{(q)} u^{(i)}(B) = c^{(q)}$$

for $q = 0, \dots, m-1$. These schemes are of the form

$$(18) \quad L_h u_h(t_k) := \sum_{i=0}^m \rho_{k,i} u_h(t_{k+i}) = L_h f(t_k) := \sum_{j=1}^J \beta_{k,j} f(\tau_{k,j})$$

$$(19) \quad M_h^{(q)} u_h := \mu_h^{(q)}[u_h] + \nu_h^{(q)}[u_h] = c^{(q)} + c_{\mu,h}^{(q)}[f] + c_{\nu,h}^{(q)}[f]$$

where $k = 0, \dots, n-m$, $q = 0, \dots, m-1$ and

$$\mu_h^{(q)}[u_h] := \sum_{i=0}^m a_i^{(q)} D_i u_h(A) + H \sum_{i=0}^{m-1} \gamma_{i,\mu}^{(q)} D_i u_h(A) H^{(i-m-1)+}$$

$$c_{\mu,h}^{(q)}[f] := H^{m-m} \sum_{j=1}^J \beta_{j,\mu}^{(q)} f(t_0 + H\xi_{j,\mu})$$

and $\nu_h^{(q)}$, $c_{\nu,h}^{(q)}$ defined analogously. Here D_i denotes the forward difference quotient

belonging to the underlying nonuniform grid

$$T_h := \{t_j, j = 0, 1, \dots, n \mid t_0 = A, t_n = B, t_{j+1} = t_j + h_j, j = 0, \dots, n-1\}$$

and h the distance $t_{m-1} - t_0$ while $(j)_+$ means the positive part of j . The meaning of the points $\tau_{k,j}, \xi_{j,u}$ is described in the papers cited.

One of the main results in [9,12] is that the difference operators can be chosen in such a way that the resulting approximations are exact for polynomials up to a certain degree and the coefficient of I_h, u_h, v_h, C_h are bounded for $h \rightarrow 0$ where the normalizing condition

$$(20) \quad \sum_{j=1}^J \beta_{k,j} = 1, k = 0, \dots, n-m$$

has been imposed. For the purposes of this section it is sufficient to assume

$$(21) \quad \max \{ |I_h p(t_k) - I_h p(t_k)|, k = 0, \dots, n-m \} \rightarrow 0 \quad (h \rightarrow 0)$$

$$(22) \quad \begin{aligned} |u_h^{(q)}[p] - u^{(q)}[p] - C_{u,h}^{(q)}[Lp]| &\rightarrow 0 \\ |v_h^{(q)}[p] - v^{(q)}[p] - C_{v,h}^{(q)}[Lp]| &\rightarrow 0 \end{aligned} \quad (h \rightarrow 0).$$

for polynomials p of degree $\leq m$ and that

$$(23) \quad \beta_{k,j}, \gamma_{i,u}^{(q)}, \gamma_{i,v}^{(q)}, \beta_{j,u}^{(q)}, \beta_{j,v}^{(q)} = O(1), h \rightarrow 0.$$

We are going to prove the following two stability inequalities.

(24) Under the conditions (20)-(23) there exists a constant $\gamma > 0$ such that for all sufficiently small h and all grid functions u_h

$$\gamma \sum_{i=0}^m \max_{k=0, \dots, n-1} |D_i u_h(t_k)| \leq \sum_{q=0}^{m-1} |D_q u_h(t_0)| + \max_{k=0, \dots, n-m} |L_h u_h(t_k)|.$$

(25) Assume (20)-(23) to hold and let (16), (17) be injective. Then there exists a constant $\gamma > 0$ such that for all sufficiently small h and all grid functions u_h

$$\gamma \sum_{i=0}^m \max_{k=0, \dots, n-1} |D_i u_h(t_k)| \leq \sum_{q=0}^{m-1} |M_h^{(q)} u_h| + \max_{k=0, \dots, n-m} |L_h u_h(t_k)|.$$

There are no restrictions made on the mesh ratios of the grids T_h . It is evident that the convergence results contained in [9,12] are an immediate consequence of (24), (25), one even obtains the slightly more general result that the m -th order difference quotients are also convergent.

In preparation for the proof of (24), (25), we rewrite the difference operator L_h in the form

$$(26) \quad L_h u_h(t_k) = \sum_{i=0}^m \alpha_{k,i} D_i u_h(t_k), \quad k = 0, \dots, n-m,$$

with certain coefficients $\alpha_{k,i}$. Inserting successively $p(t) = t^l$, $l = 0, \dots, m$, into (21), it is seen that (21) is equivalent to $(\alpha_m = 1)$

$$(27) \quad \max \{ |\alpha_{k,i} - \alpha_i(t_k)|, k = 0, \dots, n-m \} \rightarrow 0 \quad (h \rightarrow 0), \quad i = 0, \dots, m.$$

We will now apply proposition (12) to prove (25). The continuous problem (16), (17) fits into the general setting by taking

$$E = C^m[A,B], \quad F = C[A,B], \quad G^{(1)} = C^{m-1}[A,B], \quad G^{(2)} = \mathbb{R}^m$$

and defining

$$M^{(2)} : C^m[a,b] \rightarrow u + (M^{(0)}u, \dots, M^{(m-1)}u) \in \mathbb{R}^m.$$

The indexed terms are defined as discrete analogs. We write h instead of δ as index. For u_h a grid function we introduce the norms

$$\|u_h\|_l := \sum_{j=0}^l \max \{ |D_j u_h(t_k)|, k = 0, \dots, n-l \}, \quad l = m-1, m.$$

Then $E_h, G_h^{(1)}$ are taken as the vector space of grid functions on T_h equipped with the norm $\|\cdot\|_m, \|\cdot\|_{m-1}$ respectively. F_h is taken to be the vector space of grid functions w_h defined for t_k , $k = 0, \dots, n-m$, normed by

$$\|w_h\|_0 := \max \{ |w_h(t_k)|, k = 0, \dots, n-m \}.$$

Finally $G_h^{(2)} = \mathbb{R}^m$. The spaces defined so far are easily shown to form discrete approximations (see[6]). Evidently, L_h maps $E_h \rightarrow F_h$,

$$L_h^{(1)} = L_h^{(2)} := L_h.$$

$$M_h^{(2)} : E_h \rightarrow u + (M_h^{(0)}u, \dots, M_h^{(m-1)}u) \in \mathbb{R}^m.$$

We are now going to check all the conditions listed in (12). Condition (i) is proved in [6, theorem 1]. Since $K = 0$, $K_h = 0$ condition (ii) is trivially satisfied. The consistency (iii) follows from (27) and the fact that, for each $u \in E$ and $j = 0, \dots, m$,

$$\max \{ |D_j u(t_k) - u^{(j)}(t_k)|, k = 0, \dots, n-j \} \rightarrow 0 \quad (h \rightarrow 0).$$

For the same reason, the sequence $M_h^{(2)}$ is consistent with $M^{(2)}$. Moreover, due to the structure of $M_h^{(2)}$ we have

$$\|M_h^{(2)} u_h\| \leq \Gamma \|u_h\|_m, \quad u_h \in E_h,$$

with Γ independent of u_h and h , i.e. the stability of the sequence $M_h^{(2)}$.

Consistency and stability together imply (v). Condition (vi) is among the assumptions of (25). Taking (27) into account for $i = m$ it follows that $|\alpha_{k,m}| > \alpha_0 > 0$ for sufficiently small h and hence using the triangle inequality

$$\alpha_0 \|D_m u_h\|_0 \leq \|L_h u_h\|_0 + \left\| \sum_{i < m} \alpha_{\cdot, i} D_i u_h \right\|_0.$$

Because of (27), the $\alpha_{k,i}$ are uniformly bounded and we obtain for sufficiently small h

$$\alpha_0 \|u_h\|_m \leq \|L_h u_h\|_0 + \Gamma \|u_h\|_{m-1}, \quad u_h \in E_h,$$

which is the a-priori inequality (13). It remains to show (iv). Assume

$$(28) \quad J_h u_h \rightarrow g \quad (h \rightarrow 0) \quad \text{in } A(G^{(1)}, \Pi G_h^{(1)})$$

$$(29) \quad L_h u_h \rightarrow w \quad (h \rightarrow 0) \quad \text{in } A(F, \Pi F_h).$$

From (28), taking (27) into account,

$$\sum_{i < m} \alpha_{\cdot, i} D_i u_h \rightarrow \sum_{i < m} \alpha_i u^{(i)} \quad (h \rightarrow 0) \quad \text{in } A(F, \Pi F_h).$$

Then using (27) for $i = m$ and the convergence (29) we obtain

$$(30) \quad D_m u_h \rightarrow w - \sum_{i > m} \alpha_i u^{(i)} \quad (h \rightarrow 0) \quad \text{in } A(F, \Pi F_h).$$

The convergence in (28) and (30) together imply

$$u_h \rightarrow u \quad (h \rightarrow 0) \quad \text{in } A(E, \Pi E_h)$$

and hence from (30) the desired result (iv), i.e. $Lu = w$, follows. Since all assumptions of (12) have been verified, the stability inequality (1) holds, which is just (25) in our special case.

The proof of (24) is contained in the proof of (25) if the boundary conditions $M^{(q)}$ are taken to be the initial conditions specified in (24). Of course for the initial value problem the uniqueness assumption made in (25) is satisfied.

Acknowledgement. The author thanks Professor Carl deBoor for his valuable suggestions during the preparation of this report.

References

- [1] Beyn, W.J., Zur Stabilität von Differenzenverfahren für Systeme linearer gewöhnlicher Randwertaufgaben. *Numer. Math.* 29, 209-226 (1978).
- [2] Esser, H., Stabilitätsungleichungen für Diskretisierungen von Randwertaufgaben gewöhnlicher Differentialgleichungen. *Numer. Math.* 28, 69-100 (1977).
- [3] Esser, H., Stability inequalities for discrete nonlinear two-point boundary value problems. *Appl. Anal.* 10, 137-162 (1980).
- [4] Esser, H. and Niederdrenk, K., Nichtaquidistante Diskretisierungen von Randwertaufgaben. *Numer. Math.* 35, 465-478 (1980).
- [5] Grigorieff, R.D., Die Konvergenz des Rand- und Eigenwertproblems linearer gewöhnlicher Differenzengleichungen. *Numer. Math.* 15, 15-48 (1970).
- [6] Grigorieff, R.D., Zur diskreten Kompaktheit von Funktionen auf nichtaquidistanten Gittern in \mathbb{R} . *Numer. Funct. Anal. Optimiz.* 4(4), 383-395 (1981-1982).
- [7] Keller, H.B. and White, A.B., Difference methods for boundary value problems in ordinary differential equations. *SIAM J. Numer. Anal.* 12, 791-802 (1975).
- [8] Kreiss, H.O., Difference approximations for boundary and eigenvalue problems for ordinary differential equations. *Math. Comp.* 26, 605-624 (1972).
- [9] Lynch, R.E. and Rice, J.R., A high-order difference method for differential equations. *Math. Comp.* 34, 333-372 (1980).
- [10] Stummel, F., Diskrete Konvergenz linearer Operatoren I. *Math. Ann.* 190, 45-92 (1970).
- [11] Stummel, F., Approximation methods in analysis. Lecture Notes, Aarhus 1973.
- [12] Swartz, B., Compact, implicit difference schemes for a differential equation's side conditions. *Math. Comp.* 35, 733-746 (1980).
- [13] Vainikko, G., Funktionalanalysis der Diskretisierungsmethoden. Leipzig: Teubner 1976.

RDG/jgb

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2564	2. GOVT ACCESSION NO. AD-A134	3. RECIPIENT'S CATALOG NUMBER 535
4. TITLE (and Subtitle) ON PERTURBATIONS OF STABILITY INEQUALITIES WITH AN APPLICATION TO FINITE DIFFERENCE APPROXIMATIONS OF ODE'S		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) R. D. Grigorieff		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Scientific Computing
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE September 1983
		13. NUMBER OF PAGES 11
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Discrete approximation, inverse stability, ODE, BVP, finite differences, nonuniform grids		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In the framework of Stummel's discrete approximation theory, a perturbation theorem for inverse stability inequalities is proven. As an application, the inverse stability of compact finite difference schemes approximating two-point boundary value problems for linear ordinary differential equations on nonuniform grids is established.		